

UCD-05-11
gr-qc/0508071
August 2005

Horizon constraints and black hole entropy

S. CARLIP*

*Department of Physics
University of California
Davis, CA 95616
USA*

Abstract

To ask a question about a black hole in quantum gravity, one must restrict initial or boundary data to ensure that a black hole is actually present. For two-dimensional dilaton gravity, and probably a much wider class of theories as well, the imposition of a “stretched horizon” constraint alters the algebra of symmetries at the horizon, introducing a central term. Standard conformal field theory techniques can then be used to obtain the asymptotic density of states, reproducing the Bekenstein-Hawking entropy. The microscopic states responsible for black hole entropy can thus be viewed as “would-be pure gauge” states that become physical because the symmetry is altered by the requirement that a horizon exist.

To appear in *The Kerr spacetime: rotating black holes in general relativity*, edited by S. Scott, M. Visser, and D. Wiltshire (Cambridge University Press).

* email: carlip@physics.ucdavis.edu

1. Introduction

It has been more than thirty years since Bekenstein [1] and Hawking [2] first taught us that black holes are thermodynamic objects, with characteristic temperatures

$$T_H = \frac{\hbar\kappa}{2\pi c} \quad (1.1)$$

and entropies

$$S_{BH} = \frac{A}{4\hbar G}, \quad (1.2)$$

where κ is the surface gravity and A is the area of the event horizon. Extensive experience with thermodynamics in less exotic settings encourages us to believe that these quantities should reflect some kind of underlying statistical mechanics. The Bekenstein-Hawking entropy (1.2), for example, should count the number of microscopic states of the black hole. But by Wheeler’s famous dictum, “a black hole has no hair”: a classical, equilibrium black hole is determined completely by its mass, charge, and angular momentum, with no room for additional microscopic states to account for thermal behavior.

If black hole thermodynamics has a statistical mechanical origin, the relevant states must therefore be nonclassical. Indeed, they should be quantum gravitational—the Hawking temperature and Bekenstein-Hawking entropy depend upon both Planck’s constant $\hbar$ and Newton’s constant G . Thus the problem of black hole statistical mechanics is not just a technical question about some particular configurations of matter and gravitational fields; if we are lucky, it may give us new insight into the profound mysteries of quantum gravity.

This chapter will focus on an attempt to find a “universal” description of black hole statistical mechanics, one that involves quantum gravity but does not depend on fine details of any particular model of quantum gravity. This work is incomplete and tentative, and might ultimately prove to be wrong. But even if it is wrong, my hope is that we will learn something of value along the way.

2. Black hole entropy and the problem of universality

Ten years ago, the question of what microstates were responsible for black hole thermodynamics would have met with an almost unanimous answer: “We don’t know.” There were interesting ideas afloat, involving entanglement entropy [3] (the entropy coming from correlations between states inside and outside the horizon) and entropy of an “atmosphere” of external fields near the horizon [4], but we had nothing close to a complete description.

Today, by contrast, we suffer from embarrassment of riches. We have many candidates for the microscopic states of a black hole, all different, but all apparently giving the same result [5]. In particular, black hole entropy may count:

- Weakly coupled string and D-brane states [6, 7]: black holes can be constructed in semiclassical string theory as bound states of strings and higher-dimensional D-branes. For supersymmetric (BPS) configurations, thermodynamic properties computed at weak couplings are protected by symmetries as one dials couplings up to realistic values, so weakly coupled states can be counted to determine the entropy.

- Nonsingular geometries [8]: The weakly coupled D-brane excitations that account for black hole entropy in string theory may correspond to certain nonsingular, horizonless geometries; a typical black hole state would then be a “fuzzball” superposition of such states.
- States in a dual conformal field theory “at infinity” [9]: for black holes whose near-horizon geometry looks like anti-de Sitter space, AdS/CFT duality can be used to translate questions about thermodynamics to questions in a lower-dimensional dual, nongravitational conformal field theory.
- Spin network states crossing the horizon [10]: in loop quantum gravity, one can isolate a boundary field theory at the horizon and relate its states to the states of spin networks that “puncture” the horizon. The entropy apparently depends on one undetermined parameter, but once that parameter is fixed for one type of black hole, the approach yields the correct entropy for a wide range of other black holes.
- “Heavy” degrees of freedom in induced gravity [11]: as Sakharov first suggested [12], the Einstein-Hilbert action can be induced in a theory with no gravitational action by integrating out heavy matter fields. The Bekenstein-Hawking entropy can then be computed by counting these underlying massive degrees of freedom.
- No local states [13]: in the Euclidean path integral approach, black hole thermodynamics is determined by global topological features of spacetime rather than any local properties of the horizon. Perhaps no localized degrees of freedom are required to account for black hole entropy.
- Nongravitational states: Hawking’s original calculation of black hole radiation was based on quantum field theory in a fixed, classical black hole background. Perhaps the true degrees of freedom are not gravitational at all, but represent “entanglement entropy” [3] or the states of matter near the horizon [4].

None of these pictures has yet given us a complete model for black hole thermodynamics. But each can be used to count states for at least one class of black holes, and within its realm of applicability, each seems to give the correct Bekenstein-Hawking entropy (1.2). In an open field of investigation, the existence of competing explanations may be seen as a sign of health. But the existence of competing explanations that all *agree* is also, presumably, a sign that we are missing some deeper underlying structure.

This problem of universality occurs even within particular approaches to black hole statistical mechanics. In string theory, for example, one does not typically relate black hole entropy directly to horizon area. Rather, one constructs a particular semiclassical black hole as an assemblage of strings and D-branes; computes the entropy at weak coupling as a function of various charges; and then, separately, computes the horizon area as a function of those same charges. The results agree with the Bekenstein-Hawking formula (1.2), but the agreement has to be checked case by case. We may “know” that the next case will agree as well; but in a deep sense, we do not know why.

A simple approach to this problem is to note that black hole temperature and entropy can be computed semiclassically, so any quantum theory of gravity that has the right classical limit will have to give the “right” answer. But while this may be true, it does not really address the fundamental question: how is it that the classical theory enforces these restrictions on a quantum theory? In ordinary thermodynamics, we can determine entropy classically, by computing the volume of the relevant region of phase space; the correspondence principle then ensures that the quantum mechanical answer will be the same to lowest order. For black holes, no such calculation seems possible: once the mass, charge, and spin are fixed, there is no classical phase space left. Once again, we are missing a vital piece of the puzzle.

3. Symmetries and state-counting: conformal field theory and the Cardy formula

One possibility is that the missing piece is a classical symmetry of black hole spacetimes. Such a symmetry would be inherited by any quantum theory, regardless of the details of the quantization. At first sight, this explanation seems unlikely: we are not used to the idea that a symmetry can be strong enough to determine such detailed properties of a quantum theory as the density of states. In at least one instance, though, this is known to happen.

Consider a two-dimensional conformal field theory, defined initially on the complex plane with coordinates $(z, \bar{z})$. The holomorphic diffeomorphisms $z \rightarrow f(z)$, $\bar{z} \rightarrow \bar{f}(\bar{z})$ are symmetries of such a theory. Denote by L_n^{cl} and $\bar{L}_n^{cl}$ the generators of the transformations

$$z \rightarrow z + \epsilon z^{n+1}, \quad \bar{z} \rightarrow \bar{z} + \epsilon \bar{z}^{n+1}. \quad (3.1)$$

(The superscript *cl* means “classical.”) The Poisson brackets of these generators are almost uniquely determined by the symmetry: they form a Virasoro algebra,

$$\begin{aligned} \{L_m^{cl}, L_n^{cl}\} &= i(n-m)L_{m+n}^{cl} + \frac{ic^{cl}}{12}n(n^2-1)\delta_{m+n,0} \\ \{\bar{L}_m^{cl}, \bar{L}_n^{cl}\} &= i(n-m)\bar{L}_{m+n}^{cl} + \frac{ic^{cl}}{12}n(n^2-1)\delta_{m+n,0} \\ \{L_m^{cl}, \bar{L}_n^{cl}\} &= 0, \end{aligned} \quad (3.2)$$

where c^{cl} is a constant, the central charge [14].

When $c^{cl} = 0$, equation (3.2) is just a representation of the ordinary algebra of holomorphic vector fields,

$$\left[z^{m+1} \frac{d}{dz}, z^{n+1} \frac{d}{dz} \right] = (n-m) z^{m+n+1} \frac{d}{dz}. \quad (3.3)$$

Up to field redefinitions, the Virasoro algebra is the only central extension of this algebra. The central charge c commonly appears quantum mechanically, arising from operator reorderings. But can occur classically as well. There, it appears because the canonical generators are unique only up to the addition of constants; a central charge represents a

nontrivial cocycle [15], that is, a set of constants that cannot be removed by field redefinitions.

In 1986, Cardy found a remarkable property of such conformal field theories [16, 17]. Let Δ_0 be the smallest eigenvalue of L_0 in the spectrum, and define an “effective central charge”

$$c_{\text{eff}} = c - 24\Delta_0. \quad (3.4)$$

Then for large Δ , the density of states with eigenvalue Δ of L_0 has the asymptotic form

$$\rho(\Delta) \sim \exp \left\{ 2\pi \sqrt{\frac{c_{\text{eff}}\Delta}{6}} \right\} \rho(\Delta_0), \quad (3.5)$$

independent of any other details of the theory. The asymptotic behavior of the density of states is thus determined by a few features of the symmetry, the central charge c and the ground state conformal weight Δ_0 . In particular, theories with very different field contents can have exactly the same asymptotic density of states.

A careful proof of this result, using the method of steepest descents, is given in [18]. One can derive the logarithmic corrections to the entropy by the same methods [19]; indeed, by exploiting results from the theory of modular forms, one can obtain even higher order corrections [20, 21]. But although the mathematical derivation of the Cardy formula is relatively straightforward, I do not know of any good, intuitive *physical* explanation for (3.5). Standard derivations rely on a duality between high and low temperatures, which arises because of modular invariance: by interchanging cycles on a torus, one can trade a system on a circle of circumference L with inverse temperature β for a system on a circle of circumference β with inverse temperature L . Such a transformation relates states at high “energies” Δ to the ground state, leading ultimately to Cardy’s result. But it would be valuable to have a more direct understanding of why the density of states should be so strongly constrained by symmetry.

Note that upon quantization, after making the usual substitutions $\{, \} \rightarrow [,]/i\hbar$ and $L_m^{cl} \rightarrow L_m/\hbar$, the Virasoro algebra (3.2) becomes

$$\begin{aligned} [L_m, L_n] &= (m - n)L_{m+n} + \frac{c^{cl}}{12\hbar}m(m^2 - 1)\delta_{m+n,0} \\ [\bar{L}_m, \bar{L}_n] &= (m - n)\bar{L}_{m+n} + \frac{c^{cl}}{12\hbar}m(m^2 - 1)\delta_{m+n,0} \\ [L_m, \bar{L}_n] &= 0. \end{aligned} \quad (3.6)$$

A nonvanishing classical central charge c^{cl} thus contributes $c^{cl}/\hbar$ to the quantum central charge, and a classical conformal “charge” Δ^{cl} gives a quantum conformal weight $\Delta^{cl}/\hbar$. Hence the classical contribution to the entropy $\log \rho(\Delta)$ coming from (3.5) goes as $1/\hbar$, matching the behavior of the Bekenstein-Hawking entropy (1.2) and giving us a first hint that an approach of this sort might be productive.

The black holes we are interested in are not two dimensional, of course, and despite some interesting speculation [22], there is no proven higher-dimensional analog to the Cardy formula. But there is reason to hope that the two-dimensional result (3.5) might be relevant

to the near-horizon region of an arbitrary black hole. For instance, it is known that near a horizon, matter can be described by a two-dimensional conformal field theory [23, 24]. Indeed, in “tortoise coordinates,” the near-horizon metric in any dimension becomes

$$ds^2 = N^2(dt^2 - dr_*^2) + ds_\perp^2, \quad (3.7)$$

where the lapse function N goes to zero at the horizon. The Klein-Gordon equation then reduces to

$$(\square - m^2)\varphi = \frac{1}{N^2}(\partial_t^2 - \partial_{r_*}^2)\varphi + \mathcal{O}(1) = 0. \quad (3.8)$$

The mass and transverse excitations become negligible near the horizon: they are essentially red-shifted away relative to excitations in the r_* - t plane, leaving an effective two-dimensional conformal field theory at each point of the horizon. A similar reduction occurs for spinor and vector fields. Moreover, Jacobson and Kang have observed that the surface gravity and temperature of a stationary black hole are conformally invariant [25], and Medved et al. have recently shown that a generic stationary black hole metric has an approximate conformal symmetry near the horizon [26, 27].

4. The BTZ black hole

The first concrete evidence that conformal symmetry can determine black hole thermodynamics came from studying the (2+1)-dimensional black hole of Bañados, Teitelboim, and Zanelli [28–30]. A solution of the vacuum Einstein equations with a negative cosmological constant $\Lambda = -1/\ell^2$, the BTZ black hole is the (2+1)-dimensional analog of the Kerr-AdS metric. In Boyer-Lindquist-like coordinate, the BTZ metric takes the form

$$ds^2 = N^2 dt^2 - N^{-2} dr^2 - r^2 \left(d\phi + N^\phi dt \right)^2$$

$$\text{with } N = \left(-8GM + \frac{r^2}{\ell^2} + \frac{16G^2 J^2}{r^2} \right)^{1/2}, \quad N^\phi = -\frac{4GJ}{r^2}, \quad (4.1)$$

where M and J are the anti-de Sitter analogs of the ADM mass and angular momentum. As in 3+1 dimensions, the apparent singularities at $N = 0$ are merely coordinate singularities, and an analog of Kruskal-Szekeres coordinates can be found.

Like all vacuum spacetimes in 2+1 dimensions, the BTZ geometry has the peculiar feature of having constant curvature, and can be in fact expressed as a quotient of anti-de Sitter space by a discrete group of isometries. Nevertheless, it is a genuine black hole:

- It has a true event horizon at $r = r_+$ and, if $J \neq 0$, an inner Cauchy horizon at $r = r_-$, where

$$r_\pm^2 = 4GM\ell^2 \left\{ 1 \pm \left[1 - \left(\frac{J}{M\ell} \right)^2 \right]^{1/2} \right\}; \quad (4.2)$$

- it occurs as the end point of gravitational collapse of matter;




Figure 1: The Carter-Penrose diagram for a nonextremal BTZ black hole

- its Carter-Penrose diagram, figure 1, is essentially the same as that of an ordinary Kerr-AdS black hole;
- and, most important for our purposes, it exhibits standard black hole thermodynamics, with a temperature and entropy given by (1.1) and (1.2), where the horizon “area” is the circumference $A = 2\pi r_+$.

The thermodynamic character of the BTZ black hole may be verified in much the same way that it is for the Kerr black hole: by looking at quantum field theory in a BTZ background [31, 32]; by examining the Euclidean path integral [33] and the Brown-York microcanonical path integral [34]; by appealing to Wald’s Noether charge approach [30, 35]; and by investigating tunneling through the horizon [36, 37]. In 2+1 dimensions, a powerful new method is also available [38]: one can consider quantum gravitational perturbations induced by a classical scalar source, and then use detailed balance arguments to obtain thermodynamics. The quantitative agreement of all of these approaches gives us confidence that the thermal properties are real.

We now come to a deep mystery. Vacuum general relativity in 2+1 dimensions has no dynamical degrees of freedom. This is most easily seen by a simple counting argument—in the canonical formalism, the field is described by a spatial metric (three degrees of freedom per point) and its canonical momentum (three more degrees of freedom per point), but we also have three constraints that restrict initial values and three arbitrary coordinate choices, leaving $6 - 6 = 0$ dynamical degrees of freedom. The same conclusion may be reached by noting that the (2+1)-dimensional curvature tensor is algebraically determined by the Einstein tensor,

$$G^\mu{}_\nu = -\frac{1}{4}\epsilon^{\mu\pi\rho}\epsilon_{\nu\sigma\tau}R_{\pi\rho}{}^{\sigma\tau}, \quad (4.3)$$

so a spacetime that is empty apart from the presence of a cosmological constant necessarily has constant curvature. While the theory admits a few global “topological” excitations [39], there is no local dynamics. Where, then, can the degrees of freedom responsible for thermal behavior come from?

One piece of the answer, discovered independently by Strominger [40] and Birmingham, Sachs, and Sen [41], can be found by looking at boundary conditions at infinity. The conformal boundary of a (2+1)-dimensional asymptotically anti-de Sitter spacetime is a cylinder, so it is not surprising that the asymptotic symmetries of the BTZ black hole are described by a Virasoro algebra (3.2). It is a bit more surprising that this algebra has a central extension, but as Brown and Henneaux showed in 1986 [42], the classical central charge, computed from the standard ADM constraint algebra, is nonzero:

$$c = \frac{3\ell}{2G}. \quad (4.4)$$

Confirmation of this result has come from a path integral analysis [43], from investigating the constraint algebra in the Chern-Simons formalism [44], and from examining the conformal anomaly of the boundary stress-energy tensor [45, 46].

Moreover, the classical Virasoro “charges” L_0 and $\bar{L}_0$ can be computed within ordinary canonical general relativity, employing the same techniques that are used to determine the ADM mass [42]. Indeed, in this context the zero-modes of the diffeomorphisms (3.1) are simply linear combinations of time translations and rotations, and the corresponding conserved quantities are linear combinations of the ordinary ADM mass and angular momentum. For the BTZ black hole, in particular, one finds

$$\Delta = \frac{1}{16G\ell}(r_+ + r_-)^2, \quad \bar{\Delta} = \frac{1}{16G\ell}(r_+ - r_-)^2. \quad (4.5)$$

By the Cardy formula (3.5), with the added assumption that Δ_0 and $\bar{\Delta}_0$ are small, one then obtains an entropy

$$S = \log \rho \sim \frac{2\pi}{8G}(r_+ + r_-) + \frac{2\pi}{8G}(r_+ - r_-) = \frac{2\pi r_+}{4G}, \quad (4.6)$$

in precise agreement with the Bekenstein-Hawking entropy.

This argument is incomplete, of course: it tells us that the entropy is related to symmetries and boundary conditions at infinity, but does not explain the underlying quantum

degrees of freedom. I will argue later that this is a *good* feature, since it allows us to explain the “universality” described in section 2.. For the BTZ black hole, though, we can go a bit further.

As first observed by Achúcarro and Townsend [47] and subsequently extensively developed by Witten [48, 49], vacuum Einstein gravity in 2+1 dimensions with a negative cosmological constant is equivalent to a Chern-Simons gauge theory, with a gauge group $SL(2, \mathbb{R}) \times SL(2, \mathbb{R})$. On a compact manifold, a Chern-Simons theory is a “topological field theory,” described by a finite number of global degrees of freedom. On a manifold with boundary, however, boundary conditions can partially break the gauge invariance. As a consequence, field configurations that would ordinarily be considered gauge-equivalent become distinct at the boundary. In a manner reminiscent of the Goldstone mechanism [50], new “would-be pure gauge” degrees of freedom appear, providing new dynamical degrees of freedom.

For a Chern-Simons theory, the resulting induced boundary dynamics can be described by a Wess-Zumino-Witten model [51, 52]. For (2+1)-dimensional gravity, slightly stronger boundary conditions can further reduce the boundary theory to Liouville theory [53, 54], a result that may also be obtained directly in the metric formalism [55–58]. Whether the resulting degrees of freedom reproduce the Bekenstein-Hawking entropy remains an open question—for a recent review, see [59]—but Chen has found strong hints that a better understanding of Liouville theory might allow an explicit microscopic description of BTZ black hole thermodynamics [60].

5. Horizon constraints

The BTZ black hole offers a test case for the hypothesis that black hole entropy might be controlled by an underlying classical symmetry. But it is clearly not good enough. To start with, the computations described in the preceding section relied heavily on a very particular feature of (2+1)-dimensional asymptotically anti-de Sitter spacetime, the fact that the “boundary” at which asymptotic diffeomorphisms are defined is a two-dimensional timelike surface. A few other black holes have a similar character: the near-extremal black holes considered in string theory often have a near-horizon geometry that looks like that of a BTZ black hole, and two-dimensional methods can be used to obtain their entropy [40, 61]. But for the generic case, there is no reason to expect such a nice structure.

Moreover, the standard computations of BTZ black hole entropy use conformal symmetries at infinity. For a single, isolated black hole in 2+1 dimensions, this choice is probably harmless: there are no propagating degrees of freedom between the horizon and infinity, so it may not matter where we count the states. But even in 2+1 dimensions, the interpretation becomes unclear when there is more than one black hole present, or when the black hole is replaced by a “star” for which the BTZ solution is only the exterior geometry. If we wish to isolate the microscopic states of a particular black hole, it will be very difficult to do so using only symmetries at conformal infinity; we should presumably be looking near the horizon instead.

We are thus left with several hints. We should

- Look for “broken gauge invariance” to provide new degrees of freedom;
- Hope for an effective two-dimensional picture, which would allow us to use the Cardy formula;
- But look near horizon.

Before proceeding, though, we need to take a step back and ask a more general question. We want to investigate the statistical mechanics of a black hole. But how, exactly, do we tell that a black hole is present in the context of a fully quantum mechanical theory of gravity?

This question is frequently overlooked, because in the usual semiclassical approaches to black hole thermodynamics the answer is obvious: we fix a definite black hole background and then ask about quantum fields and gravitational fluctuations in that background. But in a full quantum theory of gravity, we cannot do that: there is no fixed background, the geometry is quantized, and the uncertainty principle prevents us from exactly specifying the metric. In general, it becomes difficult to tell whether a black hole is present or not. At best, we can ask a *conditional* question: “If a black hole with characteristic X is present, what is the probability of phenomenon Y ?”

There are two obvious ways to impose a suitable condition to give such a conditional probability. One, discussed in [62], is to treat the horizon as a “boundary” and impose appropriate black hole boundary conditions. The horizon is not, of course, a true boundary: a falling observer can cross a horizon without seeing anything special happen, and certainly doesn’t drop off the edge of the Universe. Nonetheless, we can ensure the presence of a black hole by specifying “boundary conditions” at a horizon. In a path integral formulation, for example, we can divide the manifold into two pieces along a hypersurface Σ and perform separate path integrals over the fields on each piece, with fields restricted at the “boundary” by the requirement that Σ be a suitable black hole horizon. This kind of split path integral has been studied in detail in 2+1 dimensions [63], where it yields the correct counting for the boundary degrees of freedom.

Alternatively, we can impose “horizon constraints” directly, either classically or in the quantum theory. We might, for example, construct an operator ϑ representing the expansion of a particular null surface, and restrict ourselves to states annihilated by ϑ . As we shall see below, such a restriction can alter the algebra of diffeomorphisms, allowing us to exploit the Cardy formula.

The “horizon as boundary” approach has been explored by a number of authors; see, for instance, [64–72]. A conformal symmetry in the r – t plane appears naturally, and one can obtain a Virasoro algebra with a central charge that leads to the correct Bekenstein-Hawking entropy. But the diffeomorphisms whose algebra yields that central charge—essentially, the diffeomorphisms that leave the lapse function invariant—are generated by vector fields that blow up at the horizon [73–75], and it is not clear whether they should be permitted in the theory. A closely related approach looks for an approximate conformal symmetry in the neighborhood of the horizon [76–79]. Again, one finds a Virasoro algebra with a central charge that apparently leads to the correct Bekenstein-Hawking entropy.




Figure 2: A spacelike “stretched horizon” Δ

The “horizon constraint” approach is much newer, and is not yet fully developed. Let us now examine it further. Suppose we wish to constrain our theory of gravity by requiring that some surface Σ be the horizon of a black hole. We must first decide exactly what we mean by a “horizon.” Demanding that Σ be a true event horizon seems impractical: an event horizon is determined globally, and requires that we know the entire future development of the spacetime. The most promising alternative is probably offered by the “isolated horizon” program [80]. An isolated horizon is essentially a null surface with vanishing expansion, with a few added technical conditions. Such a horizon shares many of the fundamental features of an event horizon [81]—in particular, it leads to standard black hole thermodynamics—and seems to do a good job of capturing the idea of a “local” horizon.

Isolated horizon constraints are constraints on the allowed data on a hypersurface, and in principle we should be able to use the well-developed apparatus of constrained Hamiltonian dynamics [82–84] to study such conditions. Unfortunately, though an isolated horizon is by definition a null surface, and we would require an approach akin to light cone quantization. Light cone quantization of gravity is difficult, and at this writing, such an analysis of horizon constraints has not been carried out.

We can much more easily impose constraints requiring the presence of a spacelike “stretched horizon” that becomes nearly null, as illustrated in figure 2. On such a hypersurface, it is possible to employ standard methods of constrained dynamics. As we shall see below, at least for the relatively simple model of two-dimensional dilaton gravity, the horizon constraints lead to a Virasoro algebra with a calculable central charge, allowing us to use the Cardy formula to obtain the correct Bekenstein-Hawking entropy.

6. Dilaton black holes and the Bekenstein-Hawking entropy

As noted in section 3., it is plausible that the thermodynamic properties of a black hole are determined by dynamics in the “ r - t plane.” As a first step, we can therefore

look at a dimensionally reduced theory. The full Einstein-Hilbert action in any dimension may be written exactly as an action for a two-dimensional theory: standard Kaluza-Klein techniques allow us to express the effect of “extra dimensions” in terms of scalar and gauge fields, albeit with an enormous gauge group [85]. Near a black hole horizon, though, most of these fields become negligible, and one obtains a simple two-dimensional theory, dilaton gravity [86, 87], with an action of the form

$$I = \frac{1}{16\pi G} \int d^2x \sqrt{-g} [AR + V(A)]. \quad (6.1)$$

Here, R is the two-dimensional scalar curvature, while A is a scalar field, the dilaton (often denoted φ). $V(A)$ is a potential whose detailed form depends on the higher-dimensional theory we started with; we will not need an exact expression. As the notation suggests, A is the transverse area in the higher-dimensional theory, and the expansion—the fractional rate of change of the transverse area along a null curve with null normal l^a —becomes

$$\vartheta = l^a \nabla_a A / A. \quad (6.2)$$

It is useful to reexpress the action (6.1) in terms of a null dyad (l^a, n^a) with $l^2 = n^2 = 0$, $l \cdot n = -1$. These determine “surface gravities” κ and $\bar{\kappa}$, defined by the conditions

$$\begin{aligned} \nabla_a l_b &= -\kappa n_a l_b - \bar{\kappa} l_a l_b \\ \nabla_a n_b &= \kappa n_a n_b + \bar{\kappa} l_a n_b. \end{aligned} \quad (6.3)$$

By an easy computation, the action (6.1) becomes

$$I = \int d^2x \left[\hat{\epsilon}^{ab} (2\kappa n_b \partial_a A - 2\bar{\kappa} l_b \partial_a A) + \sqrt{-g} V \right], \quad (6.4)$$

where $\hat{\epsilon}^{ab}$ is the two-dimension Levi-Civita density and I have adopted units such that $16\pi G = 1$. If we now define components of our dyad with respect to coordinates (u, v) ,

$$l = \sigma du + \alpha dv, \quad n = \beta du + \tau dv, \quad (6.5)$$

it is straightforward to find the Hamiltonian form of the action. Details may be found in [88]. The key feature is that the system contains three first-class constraints,

$$\begin{aligned} C_\perp &= \pi_\alpha' - \frac{1}{2} \pi_\alpha \pi_A - \tau V(A) \\ C_\parallel &= \pi_A A' - \alpha \pi_\alpha' - \tau \pi_\tau' \\ C_\pi &= \tau \pi_\tau - \alpha \pi_\alpha + 2A', \end{aligned} \quad (6.6)$$

where a dot denotes a derivative with respect to u , a prime a derivative with respect to v , and π_X is the momentum conjugate to the field X . $C_\perp$ and $C_\parallel$ are ordinary Hamiltonian and momentum constraints of general relativity, that is, the canonical versions of the generators

of diffeomorphisms; C_π is a disguised version of the generator of local Lorentz invariance, appearing because the pair $\{l, n\}$ is invariant under the boost $l \rightarrow fl, n \rightarrow f^{-1}n$.

We can now impose our “stretched horizon” constraints at the surface Σ defined by the condition $u = 0$. We first demand that Σ be “almost null,” i.e., that its normal be nearly equal to the null vector l^a . By (6.5), this requires that $\alpha = \epsilon_1 \ll 1$.

We next demand that Σ be “almost nonexpanding,” that is, that the expansion ϑ be “almost zero” on Σ . This condition is slightly more subtle, since the absolute scale of l^a is not fixed. While the restriction $\vartheta = 0$ is independent of this scale, a restriction of the form $\vartheta \ll 1$ clearly is not. Fortunately, though, the surface gravity κ scales identically under constant rescalings of l^a , so we can consistently require that $l^v \nabla_v A / \kappa A = \epsilon_2 \ll 1$. Rewriting these conditions in terms of canonical variables, we obtain two constraints:

$$\begin{aligned} K_1 &= \alpha - \epsilon_1 = 0 \\ K_2 &= A' - \frac{1}{2}\epsilon_2 A_+ \pi_A + \frac{a}{2}C_\pi = 0, \end{aligned} \tag{6.7}$$

where A_+ is the value of the dilaton at the horizon. The term proportional to C_π in K_2 is not necessary, but has been added for later convenience. By looking at a generic exact solution, one can verify that for $\epsilon_2 < 0$, these constraints do, in fact, define a spacelike stretched horizon of the type illustrated in figure 2.

K_1 and K_2 are not quite “constraints” in the ordinary sense of constrained Hamiltonian dynamics. In particular, they restrict allowable data only on the stretched horizon Σ , and cannot be imposed elsewhere. Nevertheless, they are similar enough to conventional constraints that many existing techniques can be used. In particular, note that the K_i have nontrivial brackets with the momentum and boost generators $C_\parallel$ and C_π , so these can no longer be considered generators of invariances of the constrained theory.* But we can fix this in a manner suggested by Bergmann and Komar many years ago [84]: we define new generators

$$\begin{aligned} C_\parallel &\rightarrow C_\parallel^* = C_\parallel + a_1 K_1 + a_2 K_2 \\ C_\pi &\rightarrow C_\pi^* = C_\pi + b_1 K_1 + b_2 K_2 \end{aligned} \tag{6.8}$$

with coefficients a_i and b_i chosen so that $\{C^*, K_i\} = 0$. Since the K_i vanish on admissible geometries—those for which our initial surface is a suitable stretched horizon—the constraints C^* are physically equivalent to the original C ; but they now preserve the horizon constraints as well.

We now make the crucial observation that the redefinitions (6.8) affect the Poisson brackets of the constraints. With the choice $a = -2$ in (6.7), it is not hard to check that

$$\begin{aligned} \{C_\parallel^*[\xi], C_\parallel^*[\eta]\} &= -C_\parallel^*[\xi\eta' - \eta\xi'] + \frac{1}{2}\epsilon_2 A_+ \int dv(\xi'\eta'' - \eta'\xi'') \\ \{C_\parallel^*[\xi], C_\pi^*[\eta]\} &= -C_\pi^*[\xi\eta'] \\ \{C_\pi^*[\xi], C_\pi^*[\eta]\} &= -\frac{1}{2}\epsilon_2 A_+ \int dv(\xi\eta' - \eta\xi') \end{aligned} \tag{6.9}$$

*The K_i have nontrivial brackets with $C_\perp$ as well, but that is not a problem: $C_\perp$ generates diffeomorphisms that move us off the initial surface Σ , and we should not expect the horizon constraints to be preserved.

where $C[\xi]$ means $\int dv \xi C$. The algebra (6.9) has a simple conformal field theoretical interpretation [14]: the $C_{\parallel}^*$ generate a Virasoro algebra with central charge

$$\frac{c}{48\pi} = -\frac{1}{2}\epsilon_2 A_+, \quad (6.10)$$

while C_{π}^* is an ordinary primary field of weight one.

To take advantage of the Cardy formula (3.5), the central charge (6.10) is not enough; we also need the classical Virasoro “charge” Δ . As in conventional approaches to black hole mechanics, this charge comes from the contribution of a boundary term that must be added to make the Virasoro generator $C_{\parallel}^*$ “differentiable” [89]. Under a variation of the fields, the momentum constraint (6.6) picks up a boundary term from integration by parts,

$$\delta C_{\parallel}[\xi] = \cdots + \xi \pi_A \delta A|_{v=v_+}, \quad (6.11)$$

at the boundary $v = v_+$. For Poisson brackets with $C_{\parallel}$ to be well-defined, this term must be canceled. We therefore add a boundary term to $C_{\parallel}$,

$$C_{\parallel}^{* \text{ bdry}}[\xi] = -\xi \pi_A A|_{v=v_+}, \quad (6.12)$$

which will give a nonvanishing classical contribution to Δ .

We also need a “mode expansion” to define the Fourier component L_0 , or, equivalently, a normalization for the “constant translation” ξ_0 . For a conformal field theory defined on a circle, or on a full complex plane with a natural complex coordinate, the mode expansion (3.1) is essentially unique. Here, though, it is not so obvious how to choose a coordinate z . As argued in [77], however, there is one particularly natural choice,

$$z = e^{2\pi i A/A_+}, \quad \xi_n = \frac{A_+}{2\pi A'} z^n, \quad (6.13)$$

where the normalization is chosen so that $[\xi_m, \xi_n] = i(n-m)\xi_{m+n}$.

Equation (6.12) then implies that

$$\Delta = C_{\parallel}^{* \text{ bdry}}[\xi_0] = -\frac{A_+}{2\pi A'} \pi_A A_+ = -\frac{A_+}{\pi \epsilon_2}, \quad (6.14)$$

where I have used the constraint $K_2 = 0$ to eliminate π_A in the last equality. Inserting (6.10) and (6.14) into the Cardy formula, assuming that Δ_0 is small, and restoring the factors of $16\pi G$ and $\hbar$, we obtain an entropy

$$S = \frac{2\pi}{16\pi G} \sqrt{\left(-\frac{24\pi\epsilon_2 A_+}{6\hbar}\right) \left(-\frac{A_+}{\pi\epsilon_2 \hbar}\right)} = \frac{A_+}{4\hbar G}, \quad (6.15)$$

exactly reproducing the Bekenstein-Hawking entropy (1.2).

7. What are the states?

I argued in section 2. that one of the main strengths of a symmetry-based derivation is that it is “universal,” that is, that it does not depend on the details of a quantum theory of gravity. Nevertheless, such a derivation does allow us to say *something* about the relevant states.

Standard approach to canonical gravity require that physical states satisfy the condition

$$C_{\parallel}|phys\rangle = C_{\pi}|phys\rangle = 0, \quad (7.1)$$

that is, that they be annihilated by the constraints. But the condition (7.1) is not consistent with a Virasoro algebra with nonvanishing central charge: schematically,

$$[C^*, C^*]|phys\rangle \sim C^*|phys\rangle + \text{const.}|phys\rangle \neq 0. \quad (7.2)$$

We must therefore weaken the physical state condition, for example by requiring only that

$$C_{\parallel}^{*(+)}|phys\rangle = 0 \quad (7.3)$$

where $C_{\parallel}^{*(+)}$ is the positive-frequency component of the momentum constraint.

It is well known in conformal field theory that such a loosening of the constraints leads to a collection of new “descendant” states $L_{-n}|phys\rangle$, which would be excluded by the stronger constraint (7.1). This phenomenon is closely analogous to the appearance of “boundary states” for the BTZ black hole, as discussed in section 4.. By relaxing the physical state constraints, we have allowed states that would formerly have been considered to be gauge-equivalent to differ physically, thus introducing a new set of “would-be pure gauge” states into our state-counting. It is an interesting open question whether the “Goldstone-like” description of section 4. can be extended to this setting.

8. Where do we go from here?

The “horizon constraint” program described in section 6. seems promising. But it is clear that important pieces are still missing. In particular:

- Neither the constraints (6.7) nor the mode choice (6.13) are unique. It is important to understand how sensitive the final results are to these choices. Some flexibility certainly exists—for example, it can be shown that the final expression for the black hole entropy does not depend on the parameter a in (6.7)—but a good deal of unexplored freedom remains.
- At this writing, the analysis has been completed only for the two-dimensional dilaton black hole. While it may be argued that this case captures the essential features of an arbitrary black hole, and explicit check of this claim is clearly needed.
- While the final expression (6.15) for the entropy is well-behaved, and exists in the limit that the “stretched horizon” approaches the true horizon, several intermediate

quantities—including the central charge c , the conformal weight Δ , and the vector fields ξ_n —behave badly at the horizon. Of course, the canonical approach used in section 6. itself only makes sense on a spacelike stretched horizon, so this breakdown is not necessarily a sign of an underlying problem. But it would clearly be desirable to perform a similar analysis in light cone quantization on a true horizon.

This last problem is reminiscent of the “horizon as boundary” approach of, for instance, [65], in which the relevant diffeomorphisms are generated by vector fields that blow up at the horizon. If this proves to be a general feature of conformal symmetry methods, it could be telling us something about “black hole complementarity” [90]: perhaps the Bekenstein-Hawking entropy is only well-defined for an observer who remains outside the horizon.

Beyond these particular issues, several more general questions must be answered before the conformal symmetry program can be taken too seriously. First, we need to understand much more about the coupling of horizon degrees of freedom to matter. Black hole thermodynamics is, after all, more than the Bekenstein-Hawking entropy; one must also demonstrate that any putative horizon degrees of freedom couple in a way that explains Hawking radiation. In this matter, the string theory computations clearly have the lead, reproducing not only the Hawking temperature but the correct gray body factors as well [7, 91]. But there are a few hints that similar results can be obtained from more general conformal symmetries [38, 92].

Second, if the contention that near-horizon symmetry is “universal” is correct, then such a symmetry must be present—albeit, perhaps, hidden—in other derivations of black hole thermodynamics. There are a few cases in which this is known to be true: for example, certain string theoretical derivations of near-extremal black hole thermodynamics exploit the conformal symmetry of the near-horizon BTZ geometry [40, 61], and the induced gravity approach can be related to a two-dimensional conformal symmetry [93]. But huge gaps still remain. One interesting avenue would be to explore the Euclidean version of the horizon constraint program, perhaps allowing us to relate the symmetry-derived count of states more directly to the Euclidean path integral. For the Euclidean black hole, the “stretched horizon” constraints (6.7) have a nice geometric interpretation, defining a circle in imaginary time around the horizon with a proper radius proportional to ϵ_2 . It may be that some of the choices we have made in defining the constraints become more natural in such a setting.

Third, we will eventually have to move away from “isolated horizons” to consider dynamical black holes, with horizons that grow as matter falls in and shrink as Hawking radiation carries away energy. It is only in such a setting that the horizon constraint program will be able to analyze such crucial problems as the information loss paradox and the final fate of an evaporating black hole. Whether this will eventually be possible remains to be seen.

Acknowledgments

This work was supported in part by the U.S. Department of Energy under grant DE-FG02-99ER40674.

References

- [1] J. D. Bekenstein, Phys. Rev. D7 (1973) 2333.
- [2] S. W. Hawking, Nature 248 (1974) 30.
- [3] L. Bombelli, R. K. Koul, J. Lee, and R. Sorkin, Phys. Rev. D34 (1986) 373.
- [4] G. 't Hooft, Nucl. Phys. B256 (1985) 727.
- [5] R. M. Wald, Living Rev. Rel. 4 (2001) 6.
- [6] A. Strominger and C. Vafa, Phys. Lett. B379 (1996) 99, hep-th/9601029.
- [7] A. W. Peet, Class. Quant. Grav. 15 (1998) 3291, hep-th/9712253.
- [8] S. D. Mathur, eprint hep-th/0502050.
- [9] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri, and Y. Oz, Phys. Rept. 323 (2000) 183, hep-th/9905111.
- [10] A. Ashtekar, J. Baez, A. Corichi, and K. Krasnov, Phys. Rev. Lett. 80 (1998) 904, gr-qc/9710007.
- [11] V. P. Frolov and D. V. Fursaev, Phys. Rev. D56 (1997) 2212, hep-th/9703178.
- [12] A. D. Sakharov, Sov. Phys. Dokl. 12 (1968) 1040, reprinted in Gen. Rel. Grav. 32 (2000) 365.
- [13] S. W. Hawking and C. J. Hunter, Phys. Rev. D59 (1999) 044025, hep-th/9808085.
- [14] P. Di Francesco, P. Mathieu, and D. Sénéchal, *Conformal Field Theory* (Springer, 1997).
- [15] V. Arnold, *Mathematical Methods of Classical Mechanics* (Springer, 1978), appendix 5.
- [16] J. A. Cardy, Nucl. Phys. B270 (1986) 186.
- [17] H. W. J. Blöte, J. A. Cardy, and M. P. Nightingale, Phys. Rev. Lett. 56 (1986) 742.
- [18] S. Carlip, Class. Quant. Grav. 15 (1998) 3609, hep-th/9806026.
- [19] S. Carlip, Class. Quant. Grav. 17 (2000) 4175, gr-qc/0005017.
- [20] D. Birmingham, I. Sachs, and S. Sen, Int. J. Mod. Phys. D10 (2001) 833, hep-th/0102155.
- [21] R. Dijkgraaf, J. M. Maldacena, G. W. Moore, and E. Verlinde, hep-th/0005003.
- [22] E. Verlinde, eprint hep-th/0008140.

- [23] D. Birmingham, K. S. Gupta, and S. Sen, Phys. Lett. B505 (2001) 191, hep-th/0102051.
- [24] K. S. Gupta and S. Sen, Phys. Lett. B526 (2002) 121, hep-th/0112041.
- [25] T. Jacobson and G. Kang, Class. Quant. Grav. 10 (1993) L201, gr-qc/9307002.
- [26] A. J. M. Medved, D. Martin, and M. Visser, Class. Quant. Grav. 21 (2004) 3111, gr-qc/0402069.
- [27] A. J. M. Medved, D. Martin, and M. Visser, Phys. Rev. D70 (2004) 024009, gr-qc/0403026.
- [28] M. Bañados, C. Teitelboim, and J. Zanelli, Phys. Rev. Lett. 69 (1992) 1849, hep-th/9204099.
- [29] M. Bañados, M. Henneaux, C. Teitelboim, and J. Zanelli, Phys. Rev. D48 (1993) 1506, gr-qc/9302012.
- [30] S. Carlip, Class. Quant. Grav. 12 (1995) 2853, gr-qc/9506079.
- [31] G. Lifschytz and M. Ortiz, Phys. Rev. D49 (1994) 1929, gr-qc/9310008.
- [32] S. Hyun, G. H. Lee, and J. H. Yee, Phys. Lett. B322 (1994) 182.
- [33] S. Carlip and C. Teitelboim, Phys. Rev. D51 (1995) 622, gr-qc/9405070.
- [34] J. D. Brown, J. Creighton and R. B. Mann, Phys. Rev. D50 (1994) 6394, gr-qc/9405007.
- [35] S. Carlip, J. Gegenberg, and R. B. Mann, Phys. Rev. D51 (1995) 6854, gr-qc/9410021.
- [36] F. Englert and B. Reznik, Phys. Rev. D50 (1994) 2692, gr-qc/9401010.
- [37] A. J. M. Medved, Class. Quant. Grav. 19 (2002) 589, hep-th/0110289.
- [38] R. Emparan and I. Sachs, Phys. Rev. Lett. 81 (1998) 2408, hep-th/9806122.
- [39] S. Carlip, Living Rev. Rel. 8 (2005) 1, gr-qc/0409039.
- [40] A. Strominger, JHEP 02 (1998) 009, hep-th/9712251.
- [41] D. Birmingham, I. Sachs, and S. Sen, Phys. Lett. B424 (1998) 275, hep-th/9801019.
- [42] J. D. Brown and M. Henneaux, Commun. Math. Phys. 104 (1986) 207.
- [43] H. Terashima, Phys. Lett. B499 (2001) 229, hep-th/0011010.
- [44] M. Bañados, Phys. Rev. D52 (1996) 5816, hep-th/9405171.
- [45] M. Henningson and K. Skenderis, JHEP 9807 (1998) 023, hep-th/9806087.
- [46] V. Balasubramanian and P. Kraus, Commun. Math. Phys. 208 (1999) 413, hep-th/9902121.

- [47] A. Achúcarro and P. K. Townsend, Phys. Lett. B180 (1986) 89.
- [48] E. Witten, Nucl. Phys. B311 (1988) 46.
- [49] E. Witten, Commun. Math. Phys. 121 (1989) 351.
- [50] N. Kaloper and J. Terning, personal communication.
- [51] E. Witten, Commun. Math. Phys. 92 (1984) 455.
- [52] S. Elitzur, G. W. Moore, A. Schwimmer, and N. Seiberg, Nucl. Phys. B326 (1989) 108.
- [53] O. Coussaert, M. Henneaux, and P. van Driel, Class. Quant. Grav. 12 (1995) 2961, gr-qc/9506019.
- [54] M. Rومان and Ph. Spindel, Nucl. Phys. B594 (2001) 329, hep-th/0008147.
- [55] K. Skenderis and S. N. Solodukhin, Phys. Lett. B472 (2000) 316, hep-th/9910023.
- [56] K. Bautier, F. Englert, M. Rومان, and Ph. Spindel, Phys. Lett. B479 (2000) 291, hep-th/0002156.
- [57] M. Rومان and Ph. Spindel, Class. Quant. Grav. 18 (2001) 2117, gr-qc/0011005.
- [58] S. Carlip, eprint gr-qc/0501033.
- [59] S. Carlip, eprint gr-qc/0503022.
- [60] Y.-J. Chen, Class. Quant. Grav. 21 (2004) 1153, hep-th/0310234.
- [61] K. Skenderis, Lect. Notes Phys. 541 (2000) 325, hep-th/9901050.
- [62] S. Carlip, in *Constrained Dynamics and Quantum Gravity 1996*, edited by V. de Alfaro et al., Nucl. Phys. Proc. Suppl. 57 (1997) 8, gr-qc/9702017.
- [63] E. Witten, Commun. Math. Phys. 144 (1992) 189.
- [64] S. Carlip, Phys. Rev. Lett. 82 (1999) 2828, hep-th/9812013.
- [65] S. Carlip, Class. Quant. Grav. 16 (1999) 3327, gr-qc/9906126.
- [66] D. J. Navarro, J. Navarro-Salas, and P. Navarro, Nucl. Phys. B580 (2000) 311, hep-th/9911091.
- [67] J. Jing and M.-L. Yan, Phys. Rev. D63 (2001) 024003, gr-qc/0005105.
- [68] J. M. Izquierdo, J. Navarro-Salas, and P. Navarro, Class. Quant. Grav. 19 (2002) 563, hep-th/0107132.
- [69] M.-I. Park, Nucl. Phys. B634 (2002) 339, hep-th/0111224.
- [70] S. Silva, Class. Quant. Grav 19 (2002) 3947, hep-th/0204179.

- [71] M. Cvitan, S. Pallua, and P. Prester, Phys. Lett. B555 (2003) 248, hep-th/0212029.
- [72] M. Cvitan, S. Pallua, and P. Prester, Phys. Rev. D70 (2004) 084043, hep-th/0406186.
- [73] O. Dreyer, A. Ghosh, and J. Wisniewski, Class. Quant. Grav. 18 (2001) 1929, hep-th/0101117.
- [74] J. Koga, Phys. Rev. D64 (2001) 124012, gr-qc/0107096.
- [75] N. Pinamonti and L. Vanzo, Phys. Rev. D69 (2004) 084012, hep-th/0312065.
- [76] S. N. Solodukhin, Phys. Lett. B454 (1999) 213, hep-th/9812056.
- [77] S. Carlip, Phys. Rev. Lett. 88 (2002) 241301, gr-qc/0203001.
- [78] A. Giacomini and N. Pinamonti, JHEP 0302 (2003) 014, gr-qc/0301038.
- [79] S. N. Solodukhin, Phys. Rev. Lett. 92 (2004) 061302, hep-th/0310012.
- [80] A. Ashtekar, C. Beetle, and S. Fairhurst, Class. Quant. Grav. 16 (1999) L1, gr-qc/9812065.
- [81] A. Ashtekar and B. Krishnan, Living Rev. Rel. 7 (2004) 10, gr-qc/0407042.
- [82] P. A. M. Dirac, Can. J. Math. 2 (1950) 129.
- [83] P. A. M. Dirac, Can. J. Math. 3 (1951) 1.
- [84] P. G. Bergmann and A. B. Komar, Phys. Rev. Lett. 4 (1960) 432.
- [85] J. H. Yoon, Phys. Lett. B451 (1999) 296, gr-qc/0003059.
- [86] D. Louis-Martinez and G. Kunstatter, Phys. Rev. D52 (1995) 3494, gr-qc/9503016.
- [87] D. Grumiller, W. Kummer, and D. V. Vassilevich, Phys. Rept. 369 (2002) 327, hep-th/0204253.
- [88] S. Carlip, Class. Quant. Grav. 22 (2005) 1303, hep-th/0408123.
- [89] T. Regge and C. Teitelboim, Annals Phys. 88 (1974) 286.
- [90] L. Susskind, L. Thorlacius, and J. Uglum, Phys. Rev. D48 (1993) 3743, hep-th/9306069.
- [91] J. Maldacena and A. Strominger, Phys. Rev D55 (1997) 861, hep-th/9609026.
- [92] F. Larsen, Phys. Rev. D56 (1997) 1005, hep-th/9702153.
- [93] V. P. Frolov, D. Fursaev, and A. Zelnikov, JHEP 0303 (2003) 038, hep-th/0302207.